NORTH CENTRAL FOREST EXPERIMENT STATION
Research Note NC-328
1992 FOLWELL AVE ST PAUL MN 55108 FOREST SERVICE-U.S.D.A.
1985

SEED QUALITY IN FIVE JACK PINE STANDS IN NORTH CENTRAL WISCONSIN

Richard M. Jeffers[1], *Regional Geneticist,*
Rocky Mountain Region, USDA Forest Service
Lakewood, Colorado

LIBRARY FILE COPY
INTERMOUNTAIN STATION

ABSTRACT.--Hundred-seed weights from 15 trees in each stand averaged 330 mg; 100-seed weights differed significantly among trees within stands and among stands. Germination of filled seed averaged 95 percent; germination differed significantly among trees within stands, but not among stands. Detailed seed records should be kept for all seed orchards.

KEY WORDS: *Pinus banksiana,* seed weight, seed germination, variation, seed quality.

Jack pine (*Pinus banksiana* Lamb.) exhibits rapid juvenile growth, begins to produce seed at a very young age, grows on a wide variety of sites, and produces high fiber yields. When grown for short rotations (30 to 40 years) on good soils, locally adapted jack pine may produce more wood per acre than red pine or other indigenous conifers (Rudolph and Yeatman 1982). In Canada and the Lake States it is a major source of pulpwood. During 1979, approximately 599,500 cords of jack pine pulpwood were harvested in Michigan, Minnesota, and Wisconsin (Blyth and Smith 1981). This amounted to 40 percent of all conifer pulpwood harvested in these States. Jack pine currently ranks number one among the conifers in pulpwood production and use in the Lake States.

In the Lake States jack pine is commonly reproduced through clearcutting and discing of lopped and scattered branches and it ranks third in artificial regeneration. The national forests in the region plant 2 million seedlings and direct seed 350 lbs of jack pine seed annually. Production of 1.6, 0.2, and 3.3. million seedlings in Minnesota, Wisconsin, and Michigan, respectively, in 1980-1981 attest to similar substantial planting programs on state and private lands in these States. Planting is expected to increase in coming decades in response to a realization of the need to maintain species diversity and of jack pine's great biological potential when grown under intensive culture for short rotations (Zavitkovski *et al.* 1981).

A key ingredient to successful artificial regeneration is seed quality. Much information is available on quality of jack pine seed from individual cones, trees, natural stands, plantations, provenance tests, and young seed orchards (Cecich and Rudolph 1982; Eyre and Le Barron 1944; Fowler 1965; Radsliff 1981; Rudolph 1966, 1976, 1979; Rudolph and Cecich 1979; Rudolph and Yeatman 1982; Todhunter and Polk 1981). Little information, however, is available on the variation in seed quality within and among stands from a small geographic area. This paper reports on: (1) variation in weight of filled seed, and (2) variation in germination of seed from individual trees in five jack pine stands in north central Wisconsin.

METHODS

Five jack pine stands were selected in Oneida County in north central Wisconsin on private and state lands (table 1). All five stands are located within 25 miles (40 km) of each other.

Fifteen of the best phenotypes were selected in each stand for use in studies of rooting, flowering, and seed production in seedling and clonal seed orchards. Trees

[1]*This study was conducted while Jeffers was Plant Geneticist with the North Central Forest Experiment Station, U.S. Department of Agriculture, Forest Service, Forestry Sciences Laboratory, P.O. Box 898, Rhinelander, Wisconsin 54501.*

from 228 to 425 mg, a difference of 186 percent; 22 seedlots (29 percent) had 100-seed weights of less than 300 mg; 50 seedlots (67 percent) had seed weights of 300 to 400 mg; and 3 seedlots (4 percent) had weights exceeding 400 mg. These figures compare to an average 100-seed weight of 346 mg and a range from 181 mg to 639 mg for 423 commercial seedlots reported in the U.S. Department of Agriculture Forest Service Handbook No. 450 (1974). These differences in 100-seed weight among stands and among trees within stands are statistically significant (table 4). In contrast, seed weight among the three 100-seed subsamples from each seedlot was very uniform.

The average 100-seed weight of 330 mg, and range among the 75 seedlots of 228 to 425 mg, converts to an average of 304,300 and a range of 235,300 to 438,600 full seed per kilogram of seed (table 2). Comparable figures in the U.S. Department of Agriculture Forest Service Handbook No. 450 (1974) are an average of 288,800 and a range of 156,500 to 551,200 full seed per kilogram for the 423 commercial seedlots. Thus, on the average, seed from the selected trees in this study was lighter and variation among seedlots was considerably less than reported for commercial seedlots of jack pine.

With such great variation in seed weight among individual tree seedlots, nursery managers may be tempted to sort seed according to size because uniform seed size greatly simplifies mechanical sowing. Large seed may yield larger seedlings, but many studies have shown that this superior development often is temporary. Sorting according to seed size may actually eliminate some desirable genotypes entirely due to difference in seed size among individual parent trees (Silen and Osterhaus 1979).

The great variation in seed weight among seedlots found in this study emphasizes that yield of seed from an orchard on a weight basis alone is unjustified. It is more important to know the number of seed produced by the orchard, and by each clone or family within the orchard. This kind of detailed information, therefore, necessitates maintaining detailed and accurate seed production records for all seed orchard components early in the production history of the orchard. The records will be required for several seasons, until the essential parameters have been determined including percent filled seed, seed weight of filled seed, and the germination of filled seed.

SEED GERMINATION

Total Seed

Germination of total seed (before separation of filled and empty seed) averaged 72 percent 4 days after sowing and 80 percent after 10 days (table 3). Variation in 10-day average germination among the five stands ranged from 74 percent for the Owens-Illinois stand to 90 percent for the American Legion 9 stand. Variation among individual trees ranged from 44 to 97 percent. This extreme variation in germination suggests that considerable variation existed among individual tree seedlots in percent of filled seed before separation of filled and empty seed.

Full Seed

Germination as a percent of filled seed (average of 98.1 percent filled seed) averaged 91 percent 4 days after sowing and 97 percent 10 days after sowing (table 3). Among stands average germination after 10 days ranged from 96 to 98 percent; among individual tree seedlots ranged from 86 to 100 percent. The differences among individuals within stands in germination of filled seed were significant; but, differences among stands were not significant (table 4). These results

Table 3.--*Percent germination of total and filled seed*

(In percent)

Stand	Germination of all uncleaned seed			Germination[1] of filled seed		
	Average after 4 days	Average after 10 days	Range after 10 days	Average after 4 days	Average after 10 days	Range after 10 days
AL 9	85	90	85-95	92	97	86-100
AL 11	72	84	70-94	94	98	95-100
CON	71	79	52-97	90	96	87-100
N-E	65	75	57-94	88	96	88-100
O-I	68	74	44-94	92	98	94-100
Average	72	80	62-95	91	97	90-100

[1]Percent of cleaned filled seed that germinated.

Table 4.--*Analyses of variance for 100-seed weight and percent germination*

Variable	d.f.	MS	F
HUNDRED-SEED-WEIGHT			
Stands	4	21,190	3.73**[1]
Families/stands	70	5,672	111.21**
Reps/families/stands	150	51	
GERMINATION OF FULL SEED			
Stands	4	45.0	1.51 ns[1]
Families/stands	70	29.7	5.12**
Reps/families/stands	150	5.8	

[1]Significance levels: ** = 1 percent
ns = not significant.

indicate that germination of jack pine seed is under strong genetic control and variation among individuals in germination tend to even out among jack pine stands from the same general locality.

The average of 97 percent of filled seed that germinated is comparable to the 96 to 99 percent germination of filled seed reported by Cecich and Rudolph (1982), Rudolph and Cecich (1979), and Roe (1963). Sixty-three of the 75 seedlots had better than 95 percent germination of filled seed and only three seedlots had less than 90 percent germination. For 98 percent of the seedlots only 3 to 4 percent of the filled seed, as determined from X-rays, did not germinate. This apparent lack of germination may have been due to delayed germination or partial dormancy (Rudolph and Yeatman 1982), misclassification of filled seed on the radiographs, damaged seed, or to some other, possibly genetic, cause. These excellent germination results indicate that jack pine seed that appears to be filled on radiographs are viable and, with few exceptions, will germinate.

The results of this study also show that average germination of jack pine seedlots can be increased by 15 percent or more by carefully cleaning the seed. Increases of this magnitude will more than offset the additional cost of removing empty seed. Use of good quality, improved seed, can also result in good quality seedlings.

LITERATURE CITED

Baker, William Douglas. Jack pine cone and seed quantity and quality from Gogama, Ontario. Toronto, ON: University of Toronto; 1980. 135 p. M.S. thesis.

Blyth, James E.; Smith, W. Brad. Pulpwood production in the North Central Region by County, 1979. Resour. Bull. NC-56. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station; 1981. 22 p.

Cecich, Robert A.; Rudolph, Thomas D. Time of jack pine seed maturity in Lake States provenances. Can. J. For. Res. 12(2): 368-373; 1982.

Eyre, F. H.; Le Barron, Russell K. Management of jack pine stands in the Lake States. Tech. Bull. 863. St. Paul, MN: U.S. Department of Agriculture, Forest Service, Lake States Forest Experiment Station; 1944. 66 p.

Fowler, D. P. Natural self-fertilization in three jack pines and its implications in seed orchard management. For. Sci. 11(1): 56-58; 1965.

Radsliff, Wendy A. Mother tree height and seed weight effects on a north central Minnesota population of jack pine. In: Guries, R. P., ed. Proceedings, Second North Central Tree Improvement Conference; 1981 August 5-7; Lincoln, NE. Madison, WI: University of Wisconsin-Madison; 1982: 130-139.

Roe, Eugene I. Stored seed in cones of some jack pine stands, northern Minnesota. Res. Pap. LS-1. St. Paul, MN: U.S. Department of Agriculture, Forest Service, Lake States Forest Experiment Station; 1963. 14 p.

Rudolph, Thomas D. Stimulation of earlier flowering and seed production in jack pine seedlings through greenhouse and nursery culture. In: Joint Proceedings, Second Genetics Workshop SAF and the Seventh Lake States Forest Tree Improvement Conference; 1965 October 21-23; Hickory Corners, MI. Res. Pap. NC-6. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station; 1966: 80-83.

Rudolph, Thomas D. Cone set, seed yield, seed quality, and early seedling development of S_2 generation jack pine. In: Beineke, Walter F., ed. Proceedings, Tenth Central States Forest Tree Improvement Conference; 1976 September 22-23; West Lafayette, IN. West Lafayette, IN: Purdue University; 1977. 42-60.

Rudolph, Thomas D. Seed production in the first eight years and frequency of natural selfing in a simulated jack pine seedling seed orchard. In: Proceedings, Thirteenth Lake States Forest Tree Improvement Conference; 1977 August 17-18; St. Paul, MN. Gen. Tech. Rep. NC-50. St. Paul, MN: U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station; 1979: 33-47.

Rudolph, Thomas D.; Cecich, Robert A. Provenance and plantation location variation in jack pine seed yield and quality. In: Guries, R. P., ed. Proceedings, First North Central Tree Improvement Conference; 1979 August 21-23; Madison, WI. Madison, WI: University of Wisconsin-Madison; 1979: 10-20.

Rudolph, T. D.; Yeatman, C. W. Genetics of jack pine. Res. Pap. WO-38. Washington, DC: U.S. Department of Agriculture, Forest Service; 1982. 60 p.

Silen, Roy; Osterhaus, Cary. Reduction of genetic base by sizing of bulked Douglas-fir seed lots. Tree Plant. Notes 30(1); 24-30; 1979.

Todhunter, Michael N.; Polk, R. Brooks. Seed and cone production in a clonal orchard of jack pine (*Pinus banksiana*). Can. J. For. Res. 11: 512-516; 1981.

U.S. Department of Agriculture. Schopmeyer, C. S., comp. Seeds of woody plants in the United States. Agric. Handb. 450. Washington, DC: U.S. Department of Agriculture; 1974. 883 p.

Zavitkovski, J.; Jeffers, R. M.; Nienstaedt, H.; [and others]. Biomass production of several jack pine provenances at three Lake States locations. Can. J. For. Res. 11(2): 441-447; 1981.

GPO 556-170 4/85